



Enhancement Strategies for Overcoming PCR Inhibitors

Clarissa Trogon, Phil Nhan, Clara Wang, Steven B. Lee


San Jose State University, Forensic Science Program- Justice Studies Department, MacQuarrie Hall 509, One Washington Square Hall, San Jose State University, San Jose, CA 95192-0050

Abstract

The use of forensic DNA to solve crimes is well established. Forensic evidence samples are often degraded or contain environmental contaminants, which interfere with PCR amplification (3, 10, 14, 16). There are several known inhibitors to PCR: calcium, collagen, humic acid, hematin, melanin, indigo dye, and phenol-chloroform (4, 14). These inhibitors may interfere with the cell lysis or capture of components necessary for DNA extraction, by causing DNA degradation and/or inhibiting DNA polymerase amplification of target DNA (16). Three possible mechanisms of inhibition are (1) binding of the inhibitor to the polymerase, (2) interaction of the inhibitor with the DNA, and (3) interaction with the polymerase during primer extension (14). To overcome the effect of PCR inhibitors, purification from inhibitors prior to DNA extraction, removal of inhibitors during or after DNA extraction, or relief or suppression of the effect of PCR inhibitors when performing PCR can be utilized (6). Inhibitors may cause loss of signal, peak imbalance, and/or allelic dropout (14). There are several methods that are being utilized to overcome inhibition during PCR amplification, such as adding more *Taq* and BSA (10); as well as using kits designed to overcome inhibition, such as Identifiler Plus (Applied Biosystems) and PowerPlex 16 HS (Promega).

A new method to enhance amplification includes the use of novel reagents, know as PCRboost and STRboost (Biomatrix, Inc.), that have been reported to improve PCR performance five-fold or more on challenging and difficult to amplify samples (18). **This project will explore the amplification enhancement of STRboost on low quantity and low quality DNA samples that are spiked with inhibitors.** Specifically, this project will focus on samples spiked with indigo dye and phenol chloroform. Amplification with Identifiler and Identifiler PLUS kits (Applied Biosystems) of low quantity DNA samples with and without STRboost in the presence of varying amounts of inhibitors and the use of additional *Taq* polymerase and BSA were performed. Amplification utilizing kits not already designed to overcome inhibition of inhibited, low quantity DNA samples were improved using STRboost.

Methods



Results

Identifiler Kit amplification of PhOH-CHCl₃ spiked DNA with and without PCRboost




Figure 1. Amplification results of 0.62ng DNA spiked with varying amounts (0.4uL, 0.8uL, and 1.2uL) of Phenol/Chloroform/Isoamyl 25:24:1 (PhOH-CHCl₃) with and without PCRboost B using Identifiler Kit. At 0.4uL, PCRboost recovered 3 alleles at 2 loci, and nearly 4-fold enhancement was obtained for some alleles (D21S11, D3S135, D19S433, D5S818).

0.8uL PhOH-CHCl₃ spiked DNA with and without PCRboost B




Figure 2. Electropherogram of 0.62ng with 0.8uL PhOH-CHCl₃ without PCR boost (A) and with PCRboost (B), a full profile was recovered. Note: additional peaks are due to spectral overlap/pull up.

Identifiler PLUS Kit amplification of PhOH-CHCl₃ spiked DNA with and without STRboost




Figure 3. Amplification results of 0.62ng DNA spiked with varying amounts (0.6uL, 0.9uL, and 1.2uL) of PhOH-CHCl₃ with and without STRboost using the Identifiler PLUS Kit. At 1.2uL, STRboost enhanced some alleles at 9 loci, some alleles at nearly 2-fold (D5S818 and FGA). Note: At 1.2uL using Identifiler Kit, no alleles were recovered above the 50rfu threshold even with PCRboost.

Identifiler Kit amplification of Indigo Dye spiked samples with and without PCRboost and/or "Superjuice"




Figure 4. Amplification results of 0.0625 ng DNA spiked with 0.7mM of indigo dye enhanced with and without 3xTaq/BSA ("superjuice") and/or PCRboost using the Identifiler Kit. Results indicate a combination of both enhancers improved amplification but not as well as with superjuice alone.

Identifiler PLUS Kit amplification of Indigo Dye spiked samples with and without STRboost




Figure 5. Amplification results of 0.62ng DNA spiked with 3mM Indigo Dye using the Identifiler PLUS Kit with and without STRboost. At 3mM, a combination of Identifiler PLUS additives and STRboost apparently inhibits amplification.

Conclusion

- Enhancement and recovery was achieved using STRboost and other amplification enhancers on low quality and low quantity inhibited samples using the Identifiler Kit.
- Enhancement of amplification of 0.4uL PhOH-CHCl₃ inhibited DNA was also achieved (Figure 1).
- A full profile was recovered from 0.8uL PhOH-CHCl₃ spiked DNA after the addition of PCRboost (Figure 2).
- PCRboost and STRboost amplified and enhanced the peak heights of PCR products from inhibited samples however, the enhancement was not as significant on control, uninhibited samples.
- Amplification kits (Identifiler PLUS) designed to overcome inhibition did not show significant improvement with additional enhancers, such as STRboost. In fact the amplification appeared to decrease with STRboost (Figure 3 and Figure 5).
- Using the traditional Identifiler Kit with a combination of "Superjuice" (3xTaq/BSA) and PCRboost, did show enhancement; however, not as well as Superjuice alone, indicating that too many amplification enhancers may have an inhibitory effect (Figure 3, Figure 4, and Figure 5).
- PCRboost appears to enhance the amplification of loci differentially suggesting that the enhancement may be sequence and locus dependant.

Future Steps

- Evaluate different amplification strategies by modifying the Identifiler and Identifiler PLUS kit (Applied Biosystems) and PowerPlex 16 HS (Promega) protocols to include other amplification enhancing strategies with and without STRboost.
- Conduct further research and testing to understand why STRboost may have a greater affinity for certain loci.
- Evaluate results on mixtures and non-probative DNA samples with and without STRboost.
- Evaluate DNA stored in polymers and different storage buffers and tubes (e.g. DNA stored in Samplematrix and teflon tubes) with and without PCR boost to evaluate any synergistic effects of prior storage buffers and polymers on STRboost enhancement.
- Expand collaborations to include additional labs, sample types and amplification targets (e.g. Other inhibitors, mtDNA).

Acknowledgements

This research was supported by a Howard Hughes Medical Institute Grant #52006312 to J. Soto, an NSF-REU grant #DBI-0647160 to Drs. J Soto, C Ouverney & S Lee, a CASA Small Research grant to S Lee, the BTE program of J and J and support provided by Chair, Dr. Mark Correia (Justice Studies Department), Dean, Dr. Charlie Bullock (College of Applied Sciences and Arts), and Dean, Dr. Michael Parrish (College of Science) to the Forensic Science Program.

References

- Al-Soud, W. A. & Radstrom, P. (2001). Purification and characterization of PCR-inhibitory components in blood cells. *Journal of Clinical Microbiology*, 39(2), 485-493.
- Barbieri, M., Fung, R., O'Shea, C.E., Calandro, L.M., Furtado, M.R., & Stewave, J.G. (2009). Developmental validation of the Quantifiler Duo DNA Quantification Kit for simultaneous quantification of total human and human male DNA and detection of PCR inhibitors in biological samples. *Journal of Forensic Science*, 54(2), 305-319. doi: 10.1111/j.1556-4029.2008.00951.x
- Bossett, J. (2007). PCR inhibition. *Profiles in DNA*. Promega Corporation. www.promega.com
- Brenner, M.G., Frawley, H.S., Mandi, J., Calandro, L.M., Furtado, M.R., & Stewave, J.G. (2009). Developmental validation of the PrepFiler Forensic DNA Extraction Kit for extraction of genomic DNA from biological samples. *Journal of Forensic Science*, 54(3), 599-607. doi: 10.1111/j.1556-4029.2009.01013.x
- Boonings, D.J., Pei, J., Gunn, D.C., Mei, L., Thompson, J.D., Poon, H., & Marzilli, A. (2008). An instrument for automated purification of nucleic acids from contaminated forensic samples. *National Institute of Health*, 13(1), 40-48.
- Jiang, J., Albenso, K.A., Singh, A., & Xiao, L. (2005). Development of procedures for direct extraction of cryptosporidium DNA from water concentrates and for relief of PCR inhibitors. *Applied and Environmental Microbiology*, 71(3), 1125-1141. doi:10.1285/0424-6460.71.3.1125-1141
- Kemp, B.M., Monroe, C., & Smith, D.G. (2005). Repeat silica extraction: A simple technique for the removal of PCR inhibitors from DNA extractions. *Journal of Archaeological Science*, 33, 1680-1689.
- Konieczny, M.B., Kirilova, L.L., Val, E.E. & Barnes, W.M. (2009). Mutants of taq DNA polymerase resistant to PCR inhibitors allow DNA amplification from whole blood and crude soil samples. *Nucleic Acids Research*, 37(5), 1-14. doi:10.1093/nar/gkn1029
- Kontana, E.J. & Reed, F.A. (2006). Evaluation of real-time PCR amplification efficiencies to detect PCR inhibitors. *Journal of Forensic Science*, 51(4), 795-804. doi:10.1111/j.1556-4029.2006.00182.x
- Kroeder, C.A. (1990). Relief of amplification inhibition in PCR with bovine serum albumin or T4 Gene 32 protein. *Applied and Environmental Microbiology*, 62(3), 1102-1106.
- Lantz, P.G., Mattsson, M., Wadstrom, T., & Radstrom, P. (1997). Removal of PCR inhibitors from human faecal samples through the use of an aqueous two-phase system for sample preparation prior to PCR. *Journal of Microbiological Methods*, 30(3), 159-167.
- Larkin, A. & Harrison, S.A. (1999). An improved method for STR analysis of bloodstained denim. *International Journal of Legal Medicine*, 112, 388-390.
- Malone, J.J., Chang, C.W., Lagace, R.E., Wang, D.Y., Bas, J.L., McMahon, T.P., & Hennessy, L.K. (2008). Development and validation of the AmpFISTR MiniFiler PCR Amplification Kit: A miniSTR multiplex for the analysis of degraded and/or PCR inhibited DNA. *Journal of Forensic Science*, 53(4), 838-852. doi: 10.1111/j.1556-4029.2008.00700.x
- Qin, K.L., Chung, D., & McCord, B.R. (2010). A study of PCR inhibition mechanisms using real time PCR. *Journal of Forensic Science*, 55(1), 25-33. doi: 10.1111/j.1556-4029.2009.01245.x
- Stark, P.C., Mallen, K.L., Barank, K., Rossato, R., Soto, D. & Kuske, C.R. (2005). Pre-PCR DNA quantitation of soil and sediment samples: Method development and instrument design. *Soil Biology & Biochemistry*, 32, 1101-1110.
- Wilson, I.G. (1997). Inhibition and facilitation of nucleic acid amplification. *Applied and Environmental Microbiology*, 63(10), 3741-3751.
- www.appliedbiosystems.com
- www.promega.com